

**HHS PUBLIC ACCESS**

Author manuscript

J Orthop Trauma. Author manuscript; available in PMC 2018 June 01.

Published in final edited form as:

J Orthop Trauma. 2017 June ; 31(6): e179–e185. doi:10.1097/BOT.0000000000000834.

Patterns of opioid prescribing for an orthopaedic trauma population

John Ruder, MD^{1,2}, Meghan K. Wally, MSPH^{1,2}, McKell Oliverio, BS^{1,2}, Rachel B. Seymour, PhD^{1,2}, and Joseph R. Hsu, MD^{1,2} the PRIMUM Group*

¹Department of Orthopaedic Surgery; Carolinas HealthCare System, Charlotte, NC, USA

²Carolinas Trauma Network Research Center of Excellence; Carolinas HealthCare System, Charlotte, NC, USA

Abstract

Objectives—To determine opioid prescribing practices to the orthopaedic trauma (OT) population at one level I trauma center.

Design—A retrospective study of discharge prescriptions for adult OT patients. Prescription details, injury burden, and patient demographics were abstracted for patients from initial injury through 2 month follow up.

Setting—A busy level I trauma center.

Patients/Participants—Adult OT patients admitted over a 30-day period (n=110).

Intervention—All discharge and follow-up opioid prescriptions were recorded.

Main Outcome Measurements—Morphine milligram equivalents per day, Number of opioid prescriptions, type/dose of medication prescribed

Results—135 Discharge prescriptions were written for 110 patients with orthopaedic injuries during the review period (Table 1). All patients received opioids at the time of discharge (Table 2). The MMEs prescribed at the time of discharge was 114 mg (54–300 mg) for a mean of 7.21 days (2–36.7 days). While patients with pre-injury risk factors were prescribed discharge opioids for a similar duration (7.00 days, versus 7.30 days, $p=0.81$) than those without risk factors, they were prescribed significantly more MMEs than those without (130 versus 108, $p<.05$), and were more likely to receive ER/LA opioids than those without (42.11% versus 21.98%).

Conclusions—Pain management after orthopaedic trauma continues to be opioid-centric despite involving a population at risk. Further focus on prescriber and patient education, risk evaluation

Corresponding Author: Rachel Seymour, PhD, 1320 Scott Avenue, Charlotte, NC 28204, rachel.seymour@carolinashealthcare.org, Phone: (704) 355-6969, Fax: (704) 355-8708.

* A list of contributors can be found in an Acknowledgment at the end of the article.

Conflicts of Interest: The remaining authors declare no conflicts of interest.

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention (CDC).

Presented as a poster at the Southeastern Fracture Symposium, Durham, NC, January 29, 2016.

Level of Evidence: Prognostic Level IV. See Instructions for Authors for a complete description of levels of evidence.

with mitigation, guideline development, and comprehensive pain management strategies are warranted in the orthopaedic trauma population.

Keywords

prescription narcotics; opioids; orthopaedic trauma

INTRODUCTION

Opioid-based narcotics are commonly prescribed for pain control in orthopaedic surgery patients and have a high risk for abuse and dependence.^{1,2} In 2001, Joint Commission (JCHAO) introduced mandatory pain assessment standards in all patients and cited “pain is the fifth vital sign” as an example of a strategy for some clinical settings³. Concurrently, there was a massive marketing campaign by pharmaceutical companies promoting prescription opioids. This has resulted in the U.S. developing a largely pharmaceutical approach to pain management. The result is that the US consumes the majority of the world’s prescription opioids while representing less than 5% of the world’s population.⁴ Concurrently, unintentional overdose deaths have risen sharply, largely due to an increase overdoses involving prescription opioids often obtained from a secondary non-medical source.^{5,6}

Prescription opioid use now results in more deaths than cocaine and heroin combined each year.⁷ Of the medical specialties in the U.S., orthopaedic surgeons are the third highest prescribers of opioids.⁸ Studies have evaluated the prescribing habits of orthopaedic surgeons and the effect of opioids postoperatively during various procedures.^{9,10} Patients with hip and ankle fractures treated in the Netherlands who did not receive postoperative opioids have been shown to have less postoperative pain and more satisfaction than patients with the same injuries in the U.S. who received opioids.^{9,11} During this same time period, basic science studies have negatively influenced US Orthopaedic Surgeons attitudes about Non-Steroidal Anti-Inflammatory Drugs (NSAIDS) in the setting of healing fractures, despite a lack of good clinical contraindication data and widespread use internationally.^{9,11–13} This ushered in an era of opioid monotherapy for orthopaedic injuries. On the contrary, opioids are associated with worse clinical outcomes in patients with occupational musculoskeletal disorders and patients undergoing total knee arthroplasty, reverse shoulder arthroplasty, and spine surgery.^{1,10,14,15}

Physicians and Non-physician providers who manage orthopaedic trauma patients should be particularly well-versed in risk assessment and mitigation strategies, since many in this population are at greater risk.^{16–20} Massey et al. evaluated patients with a positive toxicology screen at the time of injury and found these patients to be at an increased risk for prolonged opioid use compared to those whose toxicology screen was negative.¹⁷ In evaluating additional pre-injury risk factors, Helmerhorst et al. found patients with anxiety prior to their surgery had greater pain scores and post-injury disability compared to those without anxiety.²⁰ The study also found patients who continued to use opioids 1 to 2 months past their date of surgery had more disability and psychological distress than those who were no longer taking opioids.²⁰ Patients with pre-injury opioid use are more likely to have

multiple opioid prescribers postoperatively and continue opioids for a longer duration following surgery.^{17,18} Substance abuse is another important risk factor for subsequent abuse of prescription opioids.¹³ Moreover, rates of iatrogenically-induced addiction show wide variation and predictability in the literature^{21–27} and increase with duration of exposure.^{28–31}

Over the course of the past 20 years, Extended Release/Long Acting opioids like Oxycontin gained widespread use based on retracted claims of improved safety³², and unsubstantiated claims of improved efficacy versus short acting opioids.^{33–38} “Tamper resistant” extended release opioids were developed to curb abuse and misuse.^{32,39} It has now been demonstrated that these drugs can still be abused and remain highly addictive.^{40,41} Subsequent studies have not proven that extended release narcotics have greater efficacy than immediate release.^{33–38} Despite these data and the FDA black label warning⁴² in 2001 (updated in 2015), ER/LA opioids are still commonly prescribed. Another system implemented to identify patients at risk for opioid abuse and misuse are prescription drug monitoring programs.⁴³ Despite these efforts, the opioid epidemic continues to escalate, and a more comprehensive and efficient method to stratify and identify patients at risk for nontherapeutic opioid use is needed.⁴⁴ The purpose of this study is to determine the opioid prescribing practices in the orthopaedic trauma population at a single institution.

METHODS

Setting and Patients

A retrospective study of prescribing patterns at the time of inpatient discharge in adult orthopaedic trauma patients (including operative and nonoperative cases) admitted to a level 1 trauma center over a one-month period was conducted. Patients were discharged by either the Orthopaedic Trauma team or the Trauma Critical Care team. This audit was limited to one month, since it was conducted as a basis for a series of multi-disciplinary process improvements planned across the health system gradually over the next several months to years. All Orthopaedic Trauma discharges were conducted at the single Level 1 trauma center. The expansive health system (15 hospitals, 6 freestanding Emergency Departments (EDs), 30 Urgent Cares, and 350 clinics) on a common electronic medical record system was utilized to capture follow-up data on discharged patients. This is possible because the healthcare system maintains the dominant and growing footprint in a highly consolidated healthcare market. More importantly, the system scores highly in markers of patient loyalty like patient satisfaction and employee satisfaction^{45–47}. Within system follow-up rates are quite high in both data analytic reviews and retrospective studies^{48–51}.

Data Collection and Measures

Discharge prescription details to include prescriber, drug, dose, and duration were recorded. The same details were recorded for subsequent outpatient prescriptions up to two months following discharge^{20,52}. In addition, demographic information, including gender, age, and employment status, were abstracted from the electronic medical record into REDCap (Vanderbilt University, Nashville, TN)⁵³.

Relevant medical information abstracted included mechanism of injury, injury severity score (ISS), length of stay, and toxicology results. Daily morphine milligram equivalent (MME) dose was calculated for each prescription.⁵⁴

We established a panel of subject matter experts from a diverse background of health care and public health to conduct an evidence based review to identify objective risk factors for abuse, misuse, and diversion.³⁹ This panel included members from Public Health, Orthopaedic Trauma, Addiction Medicine, Poison Control, Pain Management, Emergency Medicine, Internal Medicine, Trauma/Critical Care, and Information Technology⁵⁵. Risk factors were further limited to those that were objectively searchable in the electronic medical record. Included risk factors were: 1) current opioid prescription with >50% remaining expected (“early refills”); 2) 2+ visits to the ED or Urgent Care with onsite treatment with opioids within the 30 days prior to the prescription; 3) 3+ prescriptions for opioids within the 30 days prior to the prescription; 4) any previous presentation for opioid overdose within the EMR; and 5) any positive screen for blood alcohol content (BAC), cocaine, or marijuana within the EMR.^{56–65} We conducted focus group style interviews with prescribers to determine appropriateness of the thresholds of the risk factors. All urgent care and ED visits, as well as pre- and post-injury prescription information, was abstracted retrospectively from the EMR. Missing data was less of a concern, since limited our variables to orders/prescriptions, diagnostic codes, and lab results that are consistently and reliably documented within the EMR. Based on these risk factors, if a patient had one or more, they were classified as “at risk”.

Data Analysis

Descriptive statistics were used to characterize the patient population and rates of opioid prescribing. T-tests were performed to determine if a difference in means existed for MME, duration of prescriptions, and number of prescriptions between patients receiving extended release vs. immediate release; as well as between “at risk” patients and patients without risk factors. Z-tests were used to detect a difference in proportion of “at risk” patients between those prescribed extended release vs. immediate release opioids. A significance level of 0.05 was used. Statistical analysis was performed using SAS.

RESULTS

135 Discharge prescriptions were written for 110 patients with orthopaedic injuries during the review period (Table 1). All patients received opioids at the time of discharge (Table 2). The MMEs prescribed at the time of discharge was 114 mg (54–300 mg) for a mean of 7.21 days (2–36.7 days). There was no difference in the mean MMEs prescribed at discharge between patients who underwent surgery and those that did not (117 mg versus 113 mg, $p=0.70$).

The most prevalent age group was 30–49 (28.2%) years. The most common mechanism of injury was a motor vehicle collision (32.7%). The mean ISS was 11.2. 53% of patients had at least one operative injury. ISS was not significantly different between patients treated operatively vs. nonoperatively. Fifty percent of the patients were unemployed at their time of injury, which is a risk factor for opioid overdose.⁶³ The mean hospital length of stay was 5.4 days (range 1–25 days).

At the time of admission, 33.6% received a toxicology screen. 16.4% of all presenting patients (almost half of those who were tested) had a positive BAC or toxicology screen for cocaine or marijuana (Table 3).

Patients who were prescribed extended-release oxycodone had a mean of 1.74 opioid prescriptions provided to them at the time of discharge, compared to 1.09 for those who did not ($p=0.53$). Patients who were prescribed extended-release oxycodone at the time of discharge were prescribed significantly more MMEs than those who did not (149 mg versus 107 mg, $p<.05$). There was no difference in the ISS of patients who received extended-release oxycodone and those who didn't (12.3 versus 12.4, $p=0.971$). These ER/LA prescriptions were for a significantly longer duration after discharge (16.3 days versus 5.18 days, $p<.05$).

Twenty-five percent of the OT patients met some criteria that indicate risk for misuse, abuse, or diversion of prescription opioids upon admission for the baseline injury (Table 4). While patients with pre-injury risk factors were prescribed discharge opioids for a similar duration (7.00 days, versus 7.30 days, $p=0.81$) than those without risk factors, they were prescribed significantly more MMEs than those without (130 versus 108, $p<.05$), and were more likely to receive ER/LA opioids than those without (42.11% versus 21.98%).

During the 2 months of follow-up, 42.7% received at least 1 additional prescription for narcotics. The most commonly prescribed opioids during follow-up visits were hydrocodone-acetaminophen (36.4%), followed by oxycodone-acetaminophen (12.7%) and other opioids (2.7%). There was a mean of 0.37 visits per patient (range 0–4 visits) in which an opioid prescription was provided during the follow-up period. Patients were provided an average of 1.1 opioid prescriptions at their follow-up visit (range 1–4 prescriptions). The mean MMEs prescribed per patient during the follow-up window was 99.2 (0–555). Of patients that received a follow-up prescription, those with pre-injury risk factors received significantly more MMEs during their follow-up than those who did not (215 versus 79.5, $p<.05$). The opioid prescriptions they received during their follow-up visits were also for a significantly longer duration (11.8 days versus 4.37 days, $p<.05$). There was no difference in the number of prescriptions provided at follow-up between patients who received extended-release oxycodone at the time of discharge and those who did not (1.17 versus 1.1, $p=0.72$). The difference in the MMEs provided at follow-up between patients who receive extended-release oxycodone at the time of discharge and those who did not was not statistically significant (142 versus 89.7, $p=0.09$). However, the opioid prescriptions they received during their follow-up visits were for a significantly longer duration (11.4 days versus 4.13 days, $p<.05$). The final opioid prescription was provided a mean of 1.82 weeks (range 0–13.3) post-discharge. Another 9.1% of patients demonstrated at-risk behavior^{56–65} during the two-month follow-up.

DISCUSSION

Until very recently, there has been very little education to prescribers on pain management strategies^{66–68} and risk other than the opioid monotherapy that has dominated the past two decades. Wide variance in prescribing practices to include the use of long-acting opioids

have been anecdotally noted in the trauma population, but publications in this area are lacking. Despite growing evidence of the link between prescription opioids and worse functional outcome, addiction, and even death, there has been a lack of translation of these data to clinical practice. Unfortunately, there has been limited education in the trauma community in the areas of risk mitigation, and non-pharmaceutical strategies for pain management.

Prescribers that are discharging orthopaedic trauma patients need to be aware that this is an at-risk population^{17,18}. Risk Evaluation and Mitigation Strategies (REMS) should be employed in these patients.^{69,70} Over 25% of the patients in this study had risk factors for abuse, misuse, or diversion of prescription opioids at the time of injury and were prescribed alarmingly high MMEs at the time of discharge. Patients who were prescribed extended-release oxycodone at the time of discharge received significantly more MMEs.

In the current study, almost 8% of this patient population had an existing/active opioid prescription with greater than 50% of the prescription remaining at the time of injury. Four-percent of the patients had received 3 or more opioid prescriptions within the 30 days prior to admission and 2% of the patients had 2 or more on-site emergency department or urgent care visits resulting in administration of opioids. Patients in this study with pre-injury risk factors received significantly more MMEs at the time of discharge, and during their follow-up visits than patients who did not. This is not only of concern because these are patients who will be at high risk of continued nontherapeutic opioid use after injury, but pre-injury opioid use has been associated with poor outcomes across multiple orthopaedic subspecialties.^{1,14,15}

In evaluating orthopaedic trauma patients at the time of admission, Massey et al. found 72% of the patients to have either a positive BAC or toxicology screen.¹⁷ This is of importance for multiple reasons. Patients with a positive toxicology screen on admission have been found to be at an increased risk for prolonged opioid use.¹⁷ Nineteen percent of the patients in this study had either a positive BAC or toxicology screen for cocaine or marijuana on admission. This group of patients is at high risk for misuse or abuse of prescription opioids and strategies to mitigate this risk should be undertaken by prescribers throughout the inpatient stay and at the time of discharge. Clinical decision support built within the electronic medical record to identify risk for abuse or misuse would be a beneficial step to engaging in primary prevention by employing multimodal strategies and cognitive behavioral therapy to appropriately reduce dose and duration of opioid prescriptions.^{71,72}

Furthermore, recognition that certain mental illnesses are risk factors for nontherapeutic opioid use⁷³ can help physicians direct more holistic treatment for these patients. Twenty-two percent of patients in this study had been diagnosed with anxiety at some point, and twenty-one percent of patients had been diagnosed with depression.

In this study of orthopaedic trauma patients who received a prescription opioid at the time of discharge, oxycodone-acetaminophen was the most commonly prescribed opioid. Almost one fifth of the patients in this study received a prescription for extended-release oxycodone at the time of discharge. The majority of these patients received an immediate release opioid

as well. Patients discharged with extended-release oxycodone averaged 149 MMEs and received 141 MMEs per day during their follow-up visits. These are both well above risk thresholds for overdose^{5,74} and the maximum of 90 MMEs per day the Centers for Disease Control and Prevention recommends for treating chronic pain.⁷⁵ Despite the attempt to reduce the danger of OxyContin® with abuse-deterrent formulations, it is now clear that this remains a highly addictive drug with abuse potential as well as potentially devastating adverse effects.^{40,41} The manufacturer of OxyContin® has paid over \$600 million in fines for inappropriately marketing the drug as non-addictive and having less potential for abuse.³² The majority of the literature citing the benefit and decreased abuse potential of extended release OxyContin® was funded by the manufacturer of the drug.^{76–79} There has been a slight slowing of the increase in opioid related deaths in some states that is thought to be a result of stricter policies and regulations rather than the reformulation of OxyContin®.^{80–83} The Academy of Orthopaedic Surgeons (AAOS) issued an information statement recommending against the use of extended-release opioids for the treatment of acute pain⁸⁴. In the current study, all of the extended-release opioid prescriptions were written by non-orthopaedic prescribers. This study highlights that despite the risks associated with extended-release opioids, they are still being prescribed at alarming rates in an at-risk population in an opioid monotherapy regimen.

In addition to opioid prescriptions at the time of discharge, 43% of patients received an additional opioid prescription during the follow-up period. Care must be taken when prescribing opioids during follow-up visits, as opioid use one to two months past the date of surgery is associated with increased disability and psychological distress.²⁰ Data suggests that 50% of patients taking prescription opioids for 3 months will continue to take them at 5 years.²⁸ During the 2 month follow-up period of this study, an additional 9% of patients developed new risk criteria.

Unfortunately, there are no current clinical practice guidelines on pain management in Orthopaedic Surgery to include Orthopaedic Trauma⁸⁵. Some guidelines do exist, like the CDC guidelines for chronic pain⁷⁵ and guidelines from other specialties and organizations⁸⁶. Acute post-injury or post-surgery guidelines for pain management are scarce and demonstrate significant research gaps^{87,88}.

There are several limitations of this study. 1. It is retrospective in nature and limited to prescriptions provided within our healthcare system. The primary focus of the study is discharge prescriptions which makes it generalizable to any busy level 1 trauma center. Due to the large size of the health system in such a consolidated market, the follow-up data on prescriptions has some validity. 2. The sample size is small and limited to patients admitted over a 30-day period and followed for two-months after injury. This audit of prescribing practices was limited to one month to define the problem and target gaps for improvement. Subsequent to the audit, our health system began a series of interventions as an iterative improvement process that is still ongoing toward safer prescribing and appropriate, comprehensive pain management. 3. Patients who were previously prescribed opioids for cancer pain were not excluded. This is a patient population that is known to be prescribed higher MMEs per day than the general population. The number of cancer patients however should be quite small.

Management of pain following orthopaedic trauma is complex, and requires that surgeons be attuned to need to balance patient safety with the need to adequately and appropriately manage pain. Our results show that twenty-five percent of orthopaedic trauma (OT) patients have risk factors for abuse, misuse, or diversion of prescription opioids at baseline injury. An additional 9% of orthopaedic trauma patients became at-risk in the follow-up time period. These data suggest that a move away from opioid monotherapy into a more comprehensive pain management strategy to include multimodal pain management, cryotherapy, cognitive behavioral therapy^{71,72} and others is needed. Orthopaedic practices and hospitals should develop standardized opioid prescribing protocols and encourage continuing medical education on opioid safety and appropriate use for physicians and non-physician providers that prescribe opioids.⁸⁹ To our knowledge, this study is the first to quantitatively evaluate the discharge prescribing practices in an at-risk orthopaedic trauma population. These data are essential as the first step in process improvement for safer opioid prescribing. Improvements should include the use of REMS, development and utilization of prescribing guidelines, and understanding of non-pharmaceutical strategies for pain management. Further, these data should compel clinicians to consider utilizing MME calculators and checking the state Prescription Drug Monitoring Database when prescribing opioids and to eliminate the use of ER/LA opioids for acute injury/pain. It also warrants a re-evaluation of the risk of Non-Steroidal Antiinflammatory Drugs versus opioids in the setting of musculoskeletal injury.

CONCLUSION

Pain management after orthopaedic trauma continues to be opioid-centric despite involving a population at risk. Further focus on prescriber and patient education, risk evaluation with mitigation, guideline development, and comprehensive pain management strategies are warranted in the orthopaedic trauma population.

Acknowledgments

PRIMUM Group (in alphabetical order):

Michael Beuhler, MD, Poison Information Center; Carolinas HealthCare System, Charlotte, NC, USA

Michael J. Bosse, MD, Department of Orthopaedic Surgery; Carolinas HealthCare System, Charlotte, NC

Emily Gerkin, Department of Orthopaedic Surgery; Carolinas HealthCare System, Charlotte, NC

Michael Gibbs, MD, Department of Emergency Medicine; Carolinas HealthCare System, Charlotte, NC, USA

Christopher Griggs, MD, Department of Emergency Medicine; Carolinas HealthCare System, Charlotte, NC, USA

Steven Jarrett, PharmD, Patient Safety; Carolinas HealthCare System, Charlotte, NC, USA

Daniel Leas, MD, Department of Orthopaedic Surgery; Carolinas HealthCare System, Charlotte, NC

Michael Runyon, MD, Department of Emergency Medicine; Carolinas HealthCare System, Charlotte, NC, USA

Animita Saha, MD, Department of Internal Medicine; Carolinas HealthCare System, Charlotte, NC, USA

Sharon Schiro, PhD, Department of Orthopaedic Surgery; Carolinas HealthCare System, Charlotte, NC

Bradley Watling, MD, Emergency Medicine Physicians; Carolinas HealthCare System, Charlotte, NC, USA

Stephen Wyatt, DO, Adult Psychiatry; Carolinas HealthCare System, Charlotte, NC, USA

Source of Funding: All authors are investigators on a grant from the Centers for Disease Control and Prevention (CE14-004 Award Number CE002520) and an internal grant from the Carolinas Trauma Network Research Center of Excellence for this work. RBS is currently receiving grants from Department of Defense, AO Foundation, and Orthopaedic Trauma Association. JRH is on the board of directors for Limb Lengthening and Reconstruction Society, is a consultant for Acumed, and is on the speaker's bureau for Smith Nephew. MJB is currently receiving a grant from the Department of Defense. MR is currently receiving a grant from NIH and royalties from UpToDate.

References

1. Zywiell MG, Stroh DA, Lee SY, Bonutti PM, Mont MA. Chronic opioid use prior to total knee arthroplasty. *J Bone Joint Surg Am.* Nov 2; 2011 93(21):1988–1993. [PubMed: 22048093]
2. Morris BJ, Mir HR. The opioid epidemic: Impact on orthopaedic surgery. *J Am Acad Orthop Surg.* 2015; 23(5):267–271. [PubMed: 25911660]
3. Joint Commission on Accreditation of Healthcare Organizations. Pain: Current Understanding of Assessment, Management, and Treatments. Washington, DC: National Pharmaceutical Council, INC and Joint Commission on Accreditation of Healthcare Organizations; 2001.
4. Manchikanti L, Singh A. Therapeutic opioids: a ten-year perspective on the complexities and complications of the escalating use, abuse, and nonmedical use of opioids. *Pain Physician.* Mar; 2008 11(2 Suppl):S63–88. [PubMed: 18443641]
5. Bohnert ASB, Valenstein M, Bair MJ, et al. Association Between Opioid Prescribing Patterns and Opioid Overdose-Related Deaths. *JAMA.* 2011; 305(13):1315–1321. [PubMed: 21467284]
6. Substance Abuse and Mental Health Services Administration. Results from the 2010 National Survey on Drug Use and Health: Summary of National Findings. U.S. Department of Health and Human Services; 2011.
7. Manchikanti L, Helm S 2nd, Fellows B, et al. Opioid epidemic in the United States. *Pain Physician.* Jul; 2012 15(3 Suppl):ES9–38. [PubMed: 22786464]
8. Volkow ND, McLellan TA, Cotto JH. Characteristics of opioid prescriptions in 2009. *JAMA.* 2011; 305(13):1299–1300. [PubMed: 21467282]
9. Lindenhovius AL, Helmerhorst GT, Schnellen AC, Vrahas M, Ring D, Kloen P. Differences in prescription of narcotic pain medication after operative treatment of hip and ankle fractures in the United States and The Netherlands. *J Trauma.* Jul; 2009 67(1):160–164. [PubMed: 19590328]
10. Kidner CL, Mayer TG, Gatchel RJ. Higher opioid doses predict poorer functional outcome in patients with chronic disabling occupational musculoskeletal disorders. *J Bone Joint Surg Am.* Apr; 2009 91(4):919–927. [PubMed: 19339577]
11. Helmerhorst GT, Lindenhovius AL, Vrahas M, Ring D, Kloen P. Satisfaction with pain relief after operative treatment of an ankle fracture. *Injury.* Nov; 2012 43(11):1958–1961. [PubMed: 22901424]
12. Kurmis AP, Kurmis TP, O'Brien JX, Dalen T. The effect of nonsteroidal anti-inflammatory drug administration on acute phase fracture-healing: a review. *J Bone Joint Surg Am.* May 2; 2012 94(9):815–823. [PubMed: 22552671]
13. Dodwell E, Latorre J, Parisini E, et al. NSAID exposure and risk of nonunion: a meta-analysis of case-control and cohort studies. *Calcif Tissue Int.* 2010; 87(3):193–202. [PubMed: 20552333]
14. Morris BJ, Laughlin MS, Elkousy HA, Gartsman GM, Edwards TB. Preoperative opioid use and outcomes after reverse shoulder arthroplasty. *J Shoulder Elbow Surg.* Jan; 2015 24(1):11–16. [PubMed: 25037063]
15. Lee D, Armaghani S, Archer KR, et al. Preoperative Opioid Use as a Predictor of Adverse Postoperative Self-Reported Outcomes in Patients Undergoing Spine Surgery. *J Bone Joint Surg Am.* Jun 4.2014 96(11):e89. [PubMed: 24897746]
16. Levy RS, Hebert CK, Munn BG, Barrack RL. Drug and alcohol use in orthopedic trauma patients: a prospective study. *J Orthop Trauma.* 1996; 10(1):21–27. [PubMed: 8926551]
17. Massey GM, Dodds HN, Roberts CS, Servoss TJ, Blondell RD. Toxicology screening in orthopedic trauma patients predicting duration of prescription opioid use. *Journal of Addictive Diseases.* 2005; 24(4):31–41. [PubMed: 16368655]

18. Morris BJ, Zumsteg JW, Archer KR, Cash B, Mir HR. Narcotic Use and Postoperative Doctor Shopping in the Orthopaedic Trauma Population. *J Bone Joint Surg Am.* Aug 6; 2014 96(15): 1257–1262. [PubMed: 25100772]
19. Holman JE, Stoddard GJ, Higgins TF. Rates of prescription opiate use before and after injury in patients with orthopaedic trauma and the risk factors for prolonged opiate use. *J Bone Joint Surg Am.* Jun 19; 2013 95(12):1075–1080. [PubMed: 23783203]
20. Helmerhorst GT, Vranceanu AM, Vrahas M, Smith M, Ring D. Risk factors for continued opioid use one to two months after surgery for musculoskeletal trauma. *J Bone Joint Surg Am.* Mar 19; 2014 96(6):495–499. [PubMed: 24647506]
21. Fishbain DA, Rosomoff H, Rosomoff R. Drug abuse, dependence, and addiction in chronic pain patients. *Clin J Pain.* 1992; 8:77–85. [PubMed: 1633386]
22. Hoffmann N, Olofsson S, Salen B, Wickstrom L. Prevalence of abuse and dependence in chronic pain patients. *Int J Addict.* 1995; 30:919–927. [PubMed: 7558484]
23. Kouyanou K, Pither C, Wessely S. Medication misuse, abuse, and chronic dependence in chronic pain patients. *J Psychosom Res.* 1997; 43:497–504. [PubMed: 9394266]
24. Chabal C, Erjavec MK, Jacobson L, Mariano A, Chaney E. Prescription opiate abuse in chronic pain patients: Clinical criteria, incidence, and predictors. *The Clinical Journal of Pain.* 1997; 13(2): 150–155. [PubMed: 9186022]
25. Katz N, Sherburne S, Beach M, et al. Behavioral monitoring and urine toxicology testing in patients receiving long-term opioid therapy. *Anesth Analg.* 2003; 97:1097–1102. [PubMed: 14500164]
26. Reid M, Engles-Horton L, Weber M, et al. Use of opioid medications for chronic non-cancer pain. *J Gen Intern Med.* 2002; 17:173–179. [PubMed: 11929502]
27. Michna E, Jamison R, Pham L, et al. Urine toxicology screening among chronic pain patients on opioid therapy: frequency and predictability of abnormal findings. *Clin J Pain.* 2007; 23:173–179. [PubMed: 17237667]
28. Martin B, Fan M, Edlund M, DeVries A, Braden J, Sullivan M. Long-term chronic opioid therapy discontinuation rates from the TROUP study. *J Gen Intern Med.* 2011; 26:1450–1457. [PubMed: 21751058]
29. Turk DC, Okifuji A. Pain terms and taxonomies. In: Fishman, S. Ballantyne, JC., Rathmell, J., editors. *Bonica's Management of Pain.* 4th. Lippincott Williams and Wilkins; 2010. p. 14–23.
30. Braden J, Fan M, Edlund M, Martin B, DeVries A, Sullivan M. Trends in use of opioids by noncancer pain type 2000–2005 among Arkansas Medicaid and HealthCore enrollees: results from the TROUP study. *J Pain.* 2008; 9(11):1026–1035. [PubMed: 18676205]
31. Korff M, Saunders K, Thomas Ray G, et al. De facto long-term opioid therapy for noncancer pain. *Clin J Pain.* 2008; 24(6):521–527. [PubMed: 18574361]
32. Meier B. In guilty plea, OxyContin maker to pay \$600 million. *The New York Times.* May 10. 2007 2007; Business Day.
33. Hale M, Fleischmann R, Salzman R, et al. Efficacy and safety of controlled release versus immediate-release oxycodone: randomized, double-blind evaluation in chronic back pain. *Clin J Pain.* 1999; 15:179–183. [PubMed: 10524470]
34. Kaplan R, Parris W, Citron M, et al. Comparison of controlled-release and immediate-release oxycodone in cancer pain. *J Clin Oncol.* 1998; 16:3230–3237. [PubMed: 9779696]
35. Staumbaugh J, Reder R, Staumbaugh M, et al. Double-blind, randomized comparison of the analgesic and pharmacokinetic profiles of controlled- and immediate-release oral oxycodone in cancer pain patients. *J Clin Pharmacol.* 2001; 41:500–506. [PubMed: 11361046]
36. Heiskanen T, Kalso E. Controlled-release oxycodone and morphine in cancer related pain. *Pain.* 1997; 73:37–45. [PubMed: 9414055]
37. Mucci-LoRusso P, Berman B, Silberstein P, et al. Controlled-release oxycodone compared with controlled-release morphine in treatment of cancer pain: a randomized, double-blind, parallel-group study. *Eur J Pain.* 1998; 2:239–249. [PubMed: 15102384]
38. Bruera E, Belzile M, Pituskin E, et al. Randomized, double-blind, cross-over trial comparing safety and efficacy of oral controlled-release oxycodone with controlled-release morphine in patients with cancer pain. *J Clin Oncol.* 1998; 16:3222–3229. [PubMed: 9779695]

39. Katz NP, Adams EH, Chilcoat H, et al. Challenges in the development of prescription opioid abuse-deterrent formulations. *The Clinical Journal of Pain*. 2007; 23(8):648–660. [PubMed: 17885342]
40. Novak S. Comment on the United States Food and Drug Administration Docket FDA-2013-D-0045. Jan.2013 2013.
41. Cicero TJ, Ellis MS. Abuse-Deterrent Formulations and the Prescription Opioid Abuse Epidemic in the United States: Lessons Learned From OxyContin. *JAMA Psychiatry*. May; 2015 72(5):424–430. [PubMed: 25760692]
42. Administration USFaD. , editor. US Food and Drug Administration. OxyContin Medication Guide. 2015.
43. Green TC, Mann MR, Bowman SE, et al. How Does Use of a Prescription Monitoring Program Change Medical Practice? *Pain Medicine*. 2012; 13:1314–1323. [PubMed: 22845339]
44. Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. Results from the 2013 National Survey on Drug Use and Health. Rockville, MD: US Department of Health and Human Services; 2013.
45. Platonova E, Kennedy K, Shewchuk R. Understanding patient satisfaction, trust, and loyalty to primary care physicians. *Med Care Res Rev*. 2008; 65(6):696–712. [PubMed: 18723450]
46. Kessler D, Mylod D. Does patient satisfaction affect patient loyalty? *Health Care Qual Assur*. 2011; 24(4):266–273.
47. Atkins P, Marshall B, Javalgi R. Happy employees lead to loyal patients. Survey of nurses and patients shows a strong link between employee satisfaction and patient loyalty. *J Health Care Mark*. 1996; 16(4):14–23.
48. Carolinas HealthCare System. Advanced Analytics. 2016. <http://www.carolinashealthcare.org/da2>. Accessed February 12, 2016
49. Lack WD, Karunakar MA, Angerame MR, et al. Type III open tibia fractures: immediate antibiotic prophylaxis minimizes infection. *J Orthop Trauma*. Jan; 2015 29(1):1–6. [PubMed: 25526095]
50. Large T, Kellam J, Bosse M, Sims S, Althausen P, Masonis J. Locked plating of supracondylar periprosthetic femur fractures. *J Arthroplasty*. 2008 Sep; 23(6 Suppl):115–120. [PubMed: 18617358]
51. Haines N, Lack WD, Seymour R, Bosse M. Defining the lower limit of a “critical bone defect” in open diaphyseal tibia fractures. *J Orthop Trauma*. 2016 May; 30(5):e158–163. [PubMed: 27101166]
52. Vranceanu A, Bachoura A, Weening A, Vrahas M, Smith R, Ring D. Psychological factors predict disability and pain intensity after skeletal trauma. *J Bone Joint Surg Am*. 2014; 96(3):e20. [PubMed: 24500592]
53. Harris P, Taylor R, Thielke R, Payne J, Gonzalez N, Conde J. Research electronic data capture (REDCap) – A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009 Apr; 42(2):377–381. [PubMed: 18929686]
54. Centers for Disease Control and Prevention. [Accessed September 13, 2016] Calculating total daily dose of opioids for safer dosage. Available at: https://www.cdc.gov/drugoverdose/pdf/calculating_total_daily_dose-a.pdf
55. Seymour R, Leas D, Wally M, Hsu J, Group P. Prescription reporting with immediate medication utilization mapping (PRIMUM): development of an alert to improve narcotic prescribing. *BMC Medical Informatics and Decision Making*. 2016; 16(111)
56. Dormuth CR, Miller TA, Huang A, Mamdani MM, Juurlink DN. Effect of a centralized prescription network on inappropriate prescriptions for opioid analgesics and benzodiazepines. *CMAJ*. Nov 6; 2012 184(16):E852–E856. [PubMed: 22949563]
57. Yang Z, Wilsey B, Bohm M, et al. Defining risk of prescription opioid overdose: pharmacy shopping and overlapping prescriptions among long-term opioid users in medicaid. *J Pain*. May; 2015 16(5):445–453. [PubMed: 25681095]
58. Mack KA, Zhang K, Paulozzi L, Jones C. Prescription practices involving opioid analgesics among Americans with Medicaid, 2010. *J Health Care Poor Underserved*. Feb; 2015 26(1):182–198. [PubMed: 25702736]

59. Grover CA, Garmel GM. How Do Emergency Physicians Interpret Prescription Narcotic History When Assessing Patients Presenting to the Emergency Department with Pain? *Perm J*. 2012; 16(4):32–36. [PubMed: 23251114]
60. Paulozzi LJ. Prescription drug overdoses: a review. *J Safety Res*. Sep; 2012 43(4):283–289. [PubMed: 23127678]
61. Paulozzi LJ, Kilbourne EM, Shah NG, et al. A history of being prescribed controlled substances and risk of drug overdose death. *Pain Medicine*. 2012; 13:87–95. [PubMed: 22026451]
62. Colburn JL, Jasinski DR, Rastegar DA. Long-term opioid therapy, aberrant behaviors, and substance misuse: Comparison of patients treated by resident and attending physicians in a general medical clinic. *Journal of Opioid Management*. 2012; 8(3):153–160. [PubMed: 22798175]
63. Johnson EM, Lanier WA, Merrill RM, et al. Unintentional prescription opioid-related overdose deaths: description of decedents by next of kin or best contact, Utah, 2008–2009. *J Gen Intern Med*. Apr; 2012 28(4):522–529. [PubMed: 23070654]
64. Jones CM, Paulozzi LJ, Mack KA. Alcohol Involvement in Opioid Pain Reliever and Benzodiazepine Drug Abuse-Related Emergency Department Visits and Drug-Related Deaths—United States, 2010. *MMWR*. 2014; 63(40):881–885. [PubMed: 25299603]
65. Sehgal N, Manchikanti L, Smith HS. Prescription Opioid Abuse in Chronic Pain: A Review of Opioid Abuse Predictors and Strategies to Curb Opioid Abuse. *Pain Physician*. 2012; 15:ES67–ES92. [PubMed: 22786463]
66. Stanek J, Renslow M, Kallianen L. The effect of an educational program on opioid prescription patterns in hand surgery: a quality improvement program. *J Hand Surg Am*. 2015; 40(2):341–346. [PubMed: 25542435]
67. Katzman J, Comerici G, Landen MG, et al. Rules and values: a coordinated regulatory and educational approach to the public health crises of chronic pain and addiction. *Am J Public Health*. 2014; 104(8):1356–1362. [PubMed: 24922121]
68. Chiauuzzi E, Trudeau K, Zacharoff K, Bond K. Identifying primary care skills and competencies in opioid risk management. *J Contin Educ health Prof*. 2011; 31(4):231–240. [PubMed: 22189986]
69. US Food and Drug Administration. [Accessed January 26, 2017] Risk Evaluation and Mitigation Strategy (REMS) for extended-release and long-acting opioids. Available at: <http://www.fda.gov/Drugs/DrugSafety/InformationbyDrugClass/ucm163647.htm>
70. Nelson L, Perrone J. Curbing the opioid epidemic in the United States: the risk evaluation and mitigation strategy (REMS). *JAMA*. 2012 Aug; 308(5):457–458. [PubMed: 22851109]
71. Vranceanu A, Barsky A, Ring D. Psychosocial aspects of disabling musculoskeletal pain. *J Bone Joint Surg Am*. 2009; 91(8):2014–2018. [PubMed: 19651964]
72. Vranceanu A, Hageman M, Strooker J, ter Meulen D, Vrahas M, Ring D. A preliminary RCT of a mind body skills based intervention addressing mood and coping strategies in patients with acute orthopaedic trauma. *Injury*. 2015; 46(4):552–557. [PubMed: 25435134]
73. Owen GT, Burton AW, Schade CM, Passik S. Urine drug testing: current recommendations and best practices. *Pain Physician*. Jul; 2012 15(3 Suppl):ES119–133. [PubMed: 22786451]
74. Dunn KM, Saunders KW, Rutter CM, et al. Opioid Prescriptions for Chronic Pain and Overdose: A Cohort Study. *Annals of Internal Medicine*. 2010; 152:85–92. [PubMed: 20083827]
75. Dowell D, Heaegerich T, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain – United States, 2016. *MMWR*. 2016; 65:1–49.
76. Coplan PM, Kale H, Sandstrom L, Landau C, Chilcoat HD. Changes in oxycodone and heroin exposures in the National Poison Data System after introduction of extended-release oxycodone with abuse-deterrent characteristics. *Pharmacoepidemiol Drug Saf*. Dec; 2013 22(12):1274–1282. [PubMed: 24123484]
77. Alexander L, Mannion RO, Weingarten B, Fanelli RJ, Stiles GL. Development and impact of prescription opioid abuse deterrent formulation technologies. *Drug Alcohol Depend*. May 1.2014 138:1–6. [PubMed: 24613631]
78. Havens JR, Leukefeld CG, DeVeaugh-Geiss AM, Coplan P, Chilcoat HD. The impact of a reformulation of extended-release oxycodone designed to deter abuse in a sample of prescription opioid abusers. *Drug Alcohol Depend*. Jun 1.2014 139:9–17. [PubMed: 24721614]

79. Sessler NE, Downing JM, Kale H, Chilcoat HD, Baumgartner TF, Coplan PM. Reductions in reported deaths following the introduction of extended-release oxycodone (OxyContin) with an abuse-deterrent formulation. *Pharmacoepidemiol Drug Saf.* Dec; 2014 23(12):1238–1246. [PubMed: 24916486]
80. Delcher C, Wagenaar AC, Goldberger BA, Cook RL, Maldonado-Molina MM. Abrupt decline in oxycodone-caused mortality after implementation of Florida's Prescription Drug Monitoring Program. *Drug Alcohol Depend.* May 1.2015 150:63–68. [PubMed: 25746236]
81. Tavernise S. Prescription Overdose Deaths in Florida Plunge After Tougher Measures, Report Says. 2014
82. Florida Health. E-FORCSE 2014–2015 Prescription Drug Monitoring Program Annual Report. Florida Health; Dec 1. 2015 2015
83. Chen L, Hedegaard H, Warner M. Drug-poisoning deaths involving opioid analgesics: United States, 1999–2011. *NCHS Data Brief.* 2014; 166
84. American Academy of Orthopaedic Surgeons. Opioid use, misuse, and abuse in orthopaedic practice. 2015
85. American Academy of Orthopaedic Surgeons. [Accessed January 26, 2017] Clinical Practice Guidelines. Available at: <http://www.aaos.org/guidelines/?ssopc=1>
86. Franklin G. Opioids for chronic noncancer pain: A position paper of the American Academy of Neurology. *Neurology.* 2014; 83:1277–1284. [PubMed: 25267983]
87. Chou R, Gordon DB, de Leon-Casasola O, et al. Management of Postoperative Pain: A Clinical Practice Guideline From the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *J Pain.* 2016; 17(2):131–157. [PubMed: 26827847]
88. Gordon D, De Leon-Casasola O, Wu C, Sluka K, Brennan T, Chou R. Research Gaps in Practice Guidelines for Acute Postoperative Pain Management in Adults: Findings from a Review of the Evidence for an American Pain Society Clinical Practice Guideline. *J Pain.* 2016; 17(2):158–166. [PubMed: 26719073]
89. Parvizi J, Miller AG, Gandhi K. Multimodal pain management after total joint arthroplasty. *J Bone Joint Surg Am.* Jun 1; 2011 93(11):1075–1084. [PubMed: 21655901]

Table 1

Demographic data for the orthopaedic trauma population.

Characteristics	N (%)
<i>Gender</i>	
Male	70 (63.6%)
Female	40 (36.4%)
<i>Age</i>	
18–29	28 (25.5%)
30–49	31 (28.2%)
50–64	22 (20.0%)
65+	24 (21.8%)
<i>Mechanism of Injury</i>	
Motor vehicle crash	36 (32.7%)
Fall <10 feet	24 (21.8%)
Motorcycle crash	16 (15.5%)
Other	10 (9.1%)
Fall >10 feet	9 (8.2%)
<i>Employment Status</i>	
Employed	53 (48.2%)
Unemployed	55 (50.0%)

Table 2

Discharge medications for the orthopaedic trauma population.

Discharge Medication	Percentage
oxycodone-acetaminophen	55.5
hydrocodone-acetaminophen	39.1
oxycodone	20.0
other opioids	3.64
benzodiazepines	2.73
morphine	1.81

Table 3

Selected Risk Characteristics of adult orthopaedic trauma population upon admission, including toxicology results upon admission and history of mental health diagnoses

Risk Factor	N (%)
Positive Toxicology Screen at admission for Alcohol, Marijuana, or Cocaine	18 (16.4%)
History of Positive Toxicology Screen in medical record for alcohol, marijuana, or cocaine	6 (5.5%)
Positive Alcohol	7 (6.4%)
Positive Opioids	6 (5.5%)
Positive Benzodiazepine	8 (7.3%)
Positive Marijuana	11 (10.0%)
Positive Anxiety	24 (21.8%)
Positive Depression	23 (20.9%)
Positive Bi-polar/schizophrenic	2 (1.82%)

Table 4

Risk Characteristics of adult orthopaedic trauma population at discharge

Baseline Risk Characteristics	N (%)
At risk baseline (met at least 1 trigger)	28 (25.5%)
Previous presentation for overdose	0 (0%)
3+ prescriptions within previous 30 days	4 (3.6%)
2+ on-site ED administration of opioids/benzodiazepines	2 (1.8%)
Existing/active prescription with >50% remaining	8 (7.3%)